

Table 12C
2005 Pediatric Nutrition Surveillance ⁽¹⁾
New York

Summary of Trends in Growth and Anemia Indicators
Children Aged < 5 Years ⁽²⁾

Summary of Trends in Growth and Anemia Indicators ⁽³⁾												
Year	Birthweight (4)			Height and Weight							Anemia Low Hb/Hct (7)	
	Low	High		Short Stature (5)	Under- weight (6)	Over- weight (6)	≥ 2 Yrs Overweight (6)					
	Number	% <2500 g	% >4000 g	Number	% <5th	% <5th	% ≥95th	Number	% 85th-<95th	% ≥95th	Number	%
2005	57,279	9.3	6.8	440,306	4.0	7.6	12.2	197,807	16.3	15.5	297,518	12.1
2004	39,975	7.6	7.8	435,810	4.2	7.0	13.0	192,069	16.8	16.6	291,919	13.3
2003	70,739	7.6	8.0	409,105	4.4	7.2	13.4	178,642	16.5	16.8	274,354	14.1
2002	67,318	7.4	8.2	418,724	4.5	7.6	13.1	175,978	16.0	16.8	281,012	14.9
2001	53,340	7.3	8.5	410,422	4.7	6.9	13.8	186,238	16.1	16.3	313,295	14.8
2000	86,940	8.8	8.3	419,669	4.5	7.0	13.6	189,836	16.2	16.2	320,455	14.4
1999	78,096	9.1	8.1	382,459	4.3	7.2	12.9	177,489	15.8	15.4	307,425	15.0
1998	93,741	9.2	8.4	448,130	4.5	7.2	12.6	207,479	15.7	14.7	365,051	14.8
1997	89,596	9.5	8.0	446,926	4.7	7.1	12.5	208,237	15.4	14.3	368,183	14.5
1996	84,519	9.3	8.4	444,700	4.9	7.0	12.8	206,651	15.4	14.5	370,406	14.3

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.



Table 12C
2005 Pediatric Nutrition Surveillance ⁽¹⁾
New York

Summary of Trends in Growth and Anemia Indicators
Children Aged < 5 Years ⁽²⁾

Summary of Trends in Growth and Anemia Indicators ⁽³⁾												
Year	Birthweight (4)			Height and Weight							Anemia Low Hb/Hct (7)	
	Low	High		Short Stature (5)	Under-weight (6)	Over-weight (6)	≥ 2 Yrs Overweight (6)					
	Number	% <2500 g	% >4000 g	Number	% <5th	% <5th	% ≥95th	Number	% 85th-<95th	% ≥95th	Number	%
1995	83,759	9.3	8.0	444,469	5.1	7.5	12.2	203,087	14.9	13.7	368,371	15.5
1994	87,220	9.5	8.2	438,633	5.4	7.6	12.4	194,146	14.8	13.8	359,006	15.5
1993	88,482	9.6	8.1	424,037	5.8	7.8	12.1	178,048	14.5	13.2	343,167	15.8
1992	85,006	9.8	8.1	384,075	6.2	8.1	12.5	152,322	14.5	13.7	307,475	16.5
1991	82,991	10.4	7.9	365,011	6.5	8.7	12.1	139,919	13.8	13.5	289,960	17.6
1990	77,756	10.2	8.1	350,273	6.6	8.7	12.2	140,224	14.1	13.1	280,382	15.6
1989	73,393	10.4	8.1	354,388	6.7	8.4	11.7	149,413	13.5	12.3	285,460	14.7

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Infants born during the reporting period included in the analysis.

(5) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(6) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

