

**Table 21C**  
**2005 Pediatric Nutrition Surveillance** (1)  
**New York**  
**Summary of Trends in Anemia by Age**  
**Children Aged < 5 Years** (2)

Page: 1  
Run Date: 1 / 11 / 07

<b>Summary of Trends in Anemia by Age</b> (3)							
Age and Year		Anemia Low Hb/Hct (4)		Age and Year		Anemia Low Hb/Hct (4)	
		Number	%			Number	%
6 - 11 Months				18 - 23 Months			
	2005	21,816	13.3		2005	39,187	12.5
	2004	15,222	17.0		2004	37,379	14.2
	2003	16,686	18.1		2003	35,080	14.4
	2002	21,960	18.9		2002	35,118	15.5
	2001	28,890	19.5		2001	39,228	14.9
	2000	33,174	20.1		2000	39,203	14.2
	1999	36,847	21.6		1999	38,553	14.2
	1998	51,899	21.4		1998	44,760	14.3
	1997	57,217	20.7		1997	43,903	13.9
	1996	60,226	19.9		1996	44,603	13.9
12 - 17 Months				24 - 35 Months			
	2005	69,715	13.4		2005	68,223	13.2
	2004	67,685	15.1		2004	67,115	14.9
	2003	61,598	16.5		2003	63,296	15.6
	2002	61,606	16.9		2002	63,229	16.2
	2001	61,007	16.9		2001	72,029	16.3
	2000	60,639	17.3		2000	72,364	15.2
	1999	50,977	18.0		1999	69,198	15.7
	1998	57,004	17.1		1998	79,369	15.4
	1997	55,258	16.5		1997	79,845	14.9
	1996	55,524	16.0		1996	80,488	15.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

\* Percentages are not calculated if < 100 records are available for analysis after exclusions.



**Table 21C**  
**2005 Pediatric Nutrition Surveillance** (1)  
**New York**  
**Summary of Trends in Anemia by Age**  
**Children Aged < 5 Years** (2)

<b>Summary of Trends in Anemia by Age</b> (3)							
Age and Year		Anemia Low Hb/Hct (4)		Age and Year		Anemia Low Hb/Hct (4)	
		Number	%			Number	%
36 - 59 Months				Total All Age Groups			
	2005	98,577	9.8		2005	297,518	12.1
	2004	104,518	10.2		2004	291,919	13.3
	2003	97,694	10.8		2003	274,354	14.1
	2002	99,099	11.6		2002	281,012	14.9
	2001	112,141	11.4		2001	313,295	14.8
	2000	115,075	10.7		2000	320,455	14.4
	1999	111,850	11.3		1999	307,425	15.0
	1998	132,019	11.1		1998	365,051	14.8
	1997	131,960	11.0		1997	368,183	14.5
	1996	129,565	10.6		1996	370,406	14.3
All Other/Unknown							
	2005	0	*				
	2004	0	*				
	2003	0	*				
	2002	0	*				
	2001	0	*				
	2000	0	*				
	1999	0	*				
	1998	0	*				
	1997	0	*				
	1996	0	*				

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

\* Percentages are not calculated if < 100 records are available for analysis after exclusions.