

Table 6B
2003 - 2005 Pediatric Nutrition Surveillance ⁽¹⁾
New York
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years ⁽²⁾

| Comparison of Growth and Anemia Indicators by County ^(3,4) | | | | | | | | | | | | | |
|--|----------------------------|------------------------|------------------------|------------------------------------|------------------------------------|-----------------------------------|----------------------|---|----------------------|--|----------------------|---------|-------------|
| County | Birthweight ⁽⁵⁾ | | | Height and Weight | | | | | | Anemia Low Hb/Hct ⁽⁸⁾ | | | |
| | Low | High | | Short Stature ⁽⁶⁾ | Under- weight ⁽⁷⁾ | Over- weight ⁽⁷⁾ | | ≥ 2 Yrs Over- weight ⁽⁷⁾ | | | | | |
| | Number | % <2500 g (Rank) | % >4000 g (Rank) | Number | % <5th (Rank) | % <5th (Rank) | % >95th (Rank) | Number | % 85th- (Rank) | % <95th (Rank) | % >95th (Rank) | Number | % (Rank) |
| Albany | 1,794 | 10.2 (56) | 7.4 (10) | 13,595 | 3.8 (8) | 7.2 (42) | 12.8 (23) | 6,288 | 17.4 (35) | 15.0 (23) | | 8,993 | 18.6 (53) |
| Allegany | 529 | 6.4 (6) | 6.8 (6) | 3,439 | 6.5 (56) | 2.1 (2) | 12.5 (22) | 1,652 | 18.7 (48) | 13.3 (8) | | 2,232 | 9.7 (12) |
| Bronx | 21,285 | 9.3 (48) | 6.7 (3) | 168,405 | 3.7 (7) | 7.7 (50) | 14.1 (36) | 76,123 | 16.2 (13) | 18.7 (57) | | 118,566 | 15.1 (46) |
| Broome | 1,740 | 7.0 (12) | 8.9 (32) | 13,150 | 6.5 (56) | 4.1 (21) | 15.3 (51) | 6,440 | 17.3 (32) | 14.2 (13) | | 8,994 | 6.8 (4) |
| Cattaraugu | 882 | 8.6 (35) | 10.7 (48) | 5,847 | 6.0 (48) | 3.0 (14) | 14.8 (47) | 2,813 | 18.6 (47) | 15.2 (27) | | 4,023 | 6.5 (3) |
| Cayuga | 649 | 9.2 (45) | 6.3 (1) | 4,851 | 4.7 (23) | 6.7 (38) | 13.6 (32) | 2,398 | 17.6 (38) | 14.8 (21) | | 2,820 | 16.2 (49) |
| Chautauqua | 1,678 | 8.1 (28) | 8.6 (27) | 11,551 | 6.0 (48) | 3.3 (16) | 11.7 (12) | 5,609 | 17.7 (40) | 13.4 (9) | | 8,037 | 11.3 (24) |
| Chemung | 421 | 8.8 (36) | 6.4 (2) | 7,724 | 5.9 (45) | 2.9 (12) | 12.4 (20) | 3,815 | 18.7 (48) | 14.3 (14) | | 5,420 | 8.0 (6) |
| Chenango | 970 | 9.2 (45) | 9.6 (40) | 7,038 | 6.4 (54) | 2.5 (7) | 14.9 (49) | 3,371 | 18.9 (51) | 16.3 (40) | | 4,738 | 19.2 (55) |
| Clinton | 622 | 7.9 (26) | 10.9 (49) | 4,100 | 5.4 (36) | 8.8 (55) | 9.9 (3) | 1,821 | 18.5 (46) | 15.4 (32) | | 2,735 | 7.5 (5) |
| Columbia | 477 | 6.9 (8) | 11.9 (54) | 3,625 | 4.3 (22) | 4.3 (23) | 14.4 (41) | 1,720 | 16.5 (18) | 15.3 (28) | | 2,543 | 11.8 (28) |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2003 - 2005 Pediatric Nutrition Surveillance ⁽¹⁾
New York
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years ⁽²⁾

| Comparison of Growth and Anemia Indicators by County ^(3,4) | | | | | | | | | | | | |
|--|----------------------------|------------------------|------------------------|------------------------------------|---------------------|------------------------------------|-----------------------------------|---|-------------------------------|--|--------|-------------|
| County | Birthweight ⁽⁵⁾ | | | Height and Weight | | | | | | Anemia Low Hb/Hct ⁽⁸⁾ | | |
| | Low | High | | Short Stature ⁽⁶⁾ | | Under- weight ⁽⁷⁾ | Over- weight ⁽⁷⁾ | ≥ 2 Yrs Over- weight ⁽⁷⁾ | | | | |
| | Number | % <2500 g (Rank) | % >4000 g (Rank) | Number | % <5th (Rank) | % <5th (Rank) | % >95th (Rank) | Number | % 85th- <95th (Rank) | % >95th (Rank) | Number | % (Rank) |
| Cortland | 624 | 8.5 (33) | 7.9 (16) | 3,890 | 5.6 (39) | 2.3 (4) | 11.6 (11) | 1,788 | 16.7 (22) | 12.7 (5) | 2,618 | 11.7 (26) |
| Delaware | 436 | 7.3 (17) | 10.1 (44) | 2,892 | 3.8 (8) | 4.7 (26) | 13.5 (31) | 1,425 | 16.2 (13) | 15.4 (32) | 1,980 | 14.2 (42) |
| Dutchess | 1,233 | 8.8 (36) | 9.9 (42) | 8,792 | 3.2 (1) | 10.4 (62) | 12.2 (19) | 3,572 | 17.0 (24) | 15.8 (36) | 5,329 | 14.1 (41) |
| Erie | 7,310 | 9.1 (42) | 8.0 (17) | 49,884 | 4.0 (15) | 7.5 (47) | 10.4 (6) | 22,485 | 15.8 (9) | 13.0 (6) | 31,654 | 10.6 (20) |
| Essex | 279 | 9.0 (39) | 8.6 (27) | 2,229 | 4.2 (19) | 7.2 (42) | 12.8 (23) | 1,081 | 19.2 (53) | 15.4 (32) | 1,412 | 9.0 (8) |
| Franklin | 498 | 7.2 (15) | 9.8 (41) | 3,550 | 5.9 (45) | 3.8 (20) | 13.1 (26) | 1,649 | 17.2 (29) | 17.8 (51) | 2,383 | 12.8 (34) |
| Fulton | 476 | 8.2 (30) | 9.2 (36) | 3,153 | 8.4 (62) | 7.8 (51) | 13.2 (28) | 1,535 | 12.3 (2) | 11.2 (2) | 2,082 | 10.4 (18) |
| Genesee | 743 | 7.7 (21) | 8.9 (32) | 5,446 | 5.4 (36) | 6.4 (36) | 14.0 (35) | 2,652 | 18.2 (43) | 16.1 (37) | 3,687 | 13.7 (36) |
| Greene | 268 | 9.7 (55) | 7.8 (15) | 1,841 | 5.3 (34) | 5.1 (27) | 13.7 (33) | 893 | 16.2 (13) | 16.6 (41) | 1,284 | 13.8 (39) |
| Hamilton | 47 | * * | * * | 251 | 5.6 (39) | 7.6 (48) | 15.5 (53) | 119 | 14.3 (4) | 16.8 (42) | 171 | 10.5 (19) |
| Herkimer | 427 | 6.1 (5) | 12.6 (58) | 4,335 | 7.7 (61) | 2.6 (9) | 13.4 (30) | 1,972 | 15.8 (9) | 14.6 (19) | 2,925 | 24.7 (62) |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2003 - 2005 Pediatric Nutrition Surveillance ⁽¹⁾
New York
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years ⁽²⁾

| Comparison of Growth and Anemia Indicators by County ^(3,4) | | | | | | | | | | | | |
|--|----------------------------|------------------------|------------------------|------------------------------------|------------------------------------|-----------------------------------|----------------------|---|----------------------|----------------------|--|-----------|
| County | Birthweight ⁽⁵⁾ | | | Height and Weight | | | | | | | Anemia Low Hb/Hct ⁽⁸⁾ | |
| | Low | High | | Short Stature ⁽⁶⁾ | Under- weight ⁽⁷⁾ | Over- weight ⁽⁷⁾ | | ≥ 2 Yrs Over- weight ⁽⁷⁾ | | | Number % (Rank) | |
| | Number | % <2500 g (Rank) | % >4000 g (Rank) | Number | % <5th (Rank) | % <5th (Rank) | % >95th (Rank) | Number | % 85th- (Rank) | % <95th (Rank) | | |
| Jefferson | 1,712 | 6.9 (8) | 12.0 (55) | 11,120 | 4.2 (19) | 7.0 (39) | 8.2 (1) | 4,990 | 11.6 (1) | 9.2 (1) | 7,264 | 11.0 (22) |
| Kings | 31,498 | 8.2 (30) | 6.7 (3) | 289,732 | 4.1 (16) | 7.9 (52) | 11.9 (14) | 131,872 | 15.5 (7) | 15.1 (24) | 199,319 | 13.7 (36) |
| Lewis | 495 | 7.3 (17) | 8.5 (24) | 3,670 | 3.9 (12) | 7.1 (40) | 10.3 (5) | 1,800 | 14.7 (6) | 14.0 (11) | 2,511 | 13.9 (40) |
| Livingston | 236 | 5.9 (4) | 10.2 (45) | 2,939 | 5.2 (30) | 5.2 (29) | 15.8 (56) | 1,383 | 19.2 (53) | 17.7 (50) | 2,012 | 15.6 (48) |
| Madison | 650 | 6.9 (8) | 12.3 (56) | 5,138 | 5.4 (36) | 2.2 (3) | 14.4 (41) | 2,401 | 17.7 (40) | 16.1 (37) | 3,463 | 11.3 (24) |
| Monroe | 4,472 | 9.4 (49) | 7.7 (13) | 35,552 | 3.5 (4) | 8.8 (55) | 11.5 (10) | 16,363 | 16.4 (17) | 14.5 (17) | 22,796 | 9.6 (11) |
| Montgomery | 540 | 9.4 (49) | 8.0 (17) | 3,848 | 7.6 (60) | 5.7 (33) | 17.0 (62) | 1,902 | 16.3 (16) | 16.2 (39) | 2,554 | 15.1 (46) |
| Nassau | 5,441 | 8.3 (32) | 6.9 (8) | 29,988 | 4.7 (23) | 9.1 (58) | 15.4 (52) | 10,955 | 17.6 (38) | 22.4 (62) | 17,920 | 12.8 (34) |
| New York | 17,040 | 7.7 (21) | 6.8 (6) | 138,602 | 3.3 (2) | 7.3 (44) | 13.0 (25) | 61,491 | 17.3 (32) | 17.2 (46) | 96,475 | 12.2 (30) |
| Niagara | 945 | 11.2 (59) | 7.7 (13) | 12,117 | 3.9 (12) | 7.3 (44) | 12.0 (15) | 5,753 | 17.2 (29) | 15.3 (28) | 7,770 | 8.1 (7) |
| Oneida | 2,278 | 9.5 (52) | 8.9 (32) | 14,952 | 5.8 (42) | 5.7 (33) | 11.3 (8) | 6,803 | 17.1 (26) | 14.3 (14) | 10,199 | 22.7 (59) |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2003 - 2005 Pediatric Nutrition Surveillance ⁽¹⁾
New York
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years ⁽²⁾

| Comparison of Growth and Anemia Indicators by County ^(3,4) | | | | | | | | | | | | |
|--|----------------------------|------------------------|------------------------|------------------------------------|------------------------------------|-----------------------------------|----------------------|---|----------------------|--|----------------------|-----------|
| County | Birthweight ⁽⁵⁾ | | | Height and Weight | | | | | | Anemia Low Hb/Hct ⁽⁸⁾ | | |
| | Low | High | | Short Stature ⁽⁶⁾ | Under- weight ⁽⁷⁾ | Over- weight ⁽⁷⁾ | | ≥ 2 Yrs Over- weight ⁽⁷⁾ | | | | |
| | Number | % <2500 g (Rank) | % >4000 g (Rank) | Number | % <5th (Rank) | % <5th (Rank) | % >95th (Rank) | Number | % 85th- (Rank) | % <95th (Rank) | % >95th (Rank) | Number |
| Onondaga | 3,828 | 9.2 (45) | 8.7 (30) | 27,574 | 4.1 (16) | 3.6 (18) | 11.7 (12) | 12,845 | 14.3 (4) | 11.3 (4) | 18,658 | 20.3 (57) |
| Ontario | 697 | 7.0 (12) | 11.3 (52) | 4,805 | 5.2 (30) | 3.7 (19) | 15.1 (50) | 2,230 | 19.3 (55) | 15.4 (32) | 3,276 | 14.7 (43) |
| Orange | 3,039 | 7.2 (15) | 8.0 (17) | 19,212 | 5.0 (27) | 8.5 (54) | 10.7 (7) | 8,387 | 16.1 (11) | 14.5 (17) | 13,300 | 11.9 (29) |
| Orleans | 371 | 4.9 (2) | 8.4 (22) | 2,470 | 4.9 (26) | 9.6 (60) | 14.5 (45) | 1,172 | 20.7 (62) | 19.0 (58) | 1,668 | 15.0 (45) |
| Oswego | 1,368 | 7.4 (20) | 10.0 (43) | 10,454 | 6.2 (53) | 2.0 (1) | 16.7 (61) | 5,179 | 18.9 (51) | 18.4 (56) | 7,334 | 9.7 (12) |
| Otsego | 429 | 8.9 (38) | 10.5 (47) | 2,961 | 5.2 (30) | 5.8 (35) | 14.3 (39) | 1,453 | 17.5 (36) | 16.8 (42) | 1,866 | 12.5 (31) |
| Putnam | 282 | 3.2 (1) | 9.2 (36) | 1,544 | 5.6 (39) | 5.2 (29) | 15.7 (55) | 598 | 16.6 (20) | 20.7 (60) | 821 | 9.0 (8) |
| Queens | 17,146 | 7.7 (21) | 6.9 (8) | 135,942 | 3.8 (8) | 8.8 (55) | 12.1 (18) | 52,841 | 16.5 (18) | 17.4 (47) | 88,512 | 9.9 (15) |
| Rensselaer | 1,325 | 9.0 (39) | 8.3 (21) | 8,542 | 3.8 (8) | 7.6 (48) | 12.0 (15) | 4,031 | 16.8 (23) | 14.1 (12) | 5,646 | 22.8 (60) |
| Richmond | 2,488 | 9.4 (49) | 7.6 (12) | 19,259 | 3.3 (2) | 7.1 (40) | 13.1 (26) | 8,018 | 17.1 (26) | 17.6 (49) | 12,425 | 10.7 (21) |
| Rockland | 2,688 | 5.7 (3) | 8.8 (31) | 18,683 | 5.9 (45) | 8.1 (53) | 10.0 (4) | 8,259 | 15.7 (8) | 13.0 (6) | 13,870 | 14.8 (44) |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2003 - 2005 Pediatric Nutrition Surveillance ⁽¹⁾
New York
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years ⁽²⁾

| Comparison of Growth and Anemia Indicators by County ^(3,4) | | | | | | | | | | | | | |
|--|----------------------------|------------------------|------------------------|------------------------------------|------------------------------------|-----------------------------------|----------------------|---|----------------------|--|-----------------------------|--------|----------------------|
| County | Birthweight ⁽⁵⁾ | | | Height and Weight | | | | | | Anemia Low Hb/Hct ⁽⁸⁾ | | | |
| | Low | High | | Short Stature ⁽⁶⁾ | Under- weight ⁽⁷⁾ | Over- weight ⁽⁷⁾ | | ≥ 2 Yrs Over- weight ⁽⁷⁾ | | | Number % (Rank) | | |
| | Number | % <2500 g (Rank) | % >4000 g (Rank) | Number | % <5th (Rank) | % <5th (Rank) | % >95th (Rank) | Number | % 85th- (Rank) | % <95th (Rank) | | | % >95th (Rank) |
| Saratoga | 722 | 9.1 (42) | 12.3 (56) | 5,194 | 5.2 (30) | 7.3 (44) | 12.4 (20) | 2,311 | 17.0 (24) | 13.4 (9) | | 3,488 | 19.2 (55) |
| Schenectad | 998 | 9.6 (53) | 6.7 (3) | 8,311 | 4.1 (16) | 6.5 (37) | 14.3 (39) | 3,713 | 16.1 (11) | 15.1 (24) | | 5,638 | 24.5 (61) |
| Schoharie | 118 | 10.2 (56) | 9.3 (38) | 1,879 | 5.1 (29) | 4.4 (24) | 16.3 (59) | 913 | 17.5 (36) | 15.3 (28) | | 1,282 | 17.8 (52) |
| Schuyler | 54 | * * | * * | 1,110 | 6.7 (58) | 2.5 (7) | 15.8 (56) | 520 | 18.1 (42) | 19.0 (58) | | 753 | 6.0 (2) |
| Seneca | 198 | 10.6 (58) | 11.1 (51) | 1,451 | 5.8 (42) | 3.2 (15) | 14.4 (41) | 685 | 20.3 (61) | 18.0 (52) | | 996 | 20.5 (58) |
| St. Lawren | 657 | 6.7 (7) | 8.4 (22) | 4,307 | 5.8 (42) | 2.6 (9) | 14.6 (46) | 1,959 | 18.8 (50) | 18.0 (52) | | 3,014 | 18.6 (53) |
| Steuben | 1,041 | 8.0 (27) | 8.5 (24) | 7,285 | 6.7 (58) | 2.4 (5) | 13.7 (33) | 3,560 | 18.3 (44) | 14.6 (19) | | 5,026 | 11.1 (23) |
| Suffolk | 8,489 | 6.9 (8) | 8.2 (20) | 39,043 | 3.9 (12) | 9.4 (59) | 14.4 (41) | 12,398 | 19.5 (59) | 22.2 (61) | | 21,336 | 10.1 (16) |
| Sullivan | 779 | 8.1 (28) | 8.6 (27) | 4,981 | 6.4 (54) | 2.9 (12) | 14.2 (38) | 2,223 | 17.3 (32) | 17.5 (48) | | 2,948 | 10.3 (17) |
| Tioga | 521 | 7.7 (21) | 10.9 (49) | 3,994 | 6.0 (48) | 2.6 (9) | 14.8 (47) | 2,041 | 18.4 (45) | 15.1 (24) | | 2,710 | 17.7 (51) |
| Tompkins | 736 | 7.1 (14) | 9.5 (39) | 4,450 | 4.2 (19) | 5.3 (32) | 8.7 (2) | 2,036 | 13.4 (3) | 11.2 (2) | | 2,955 | 12.6 (32) |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2003 - 2005 Pediatric Nutrition Surveillance ⁽¹⁾
New York
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years ⁽²⁾

| Comparison of Growth and Anemia Indicators by County ^(3,4) | | | | | | | | | | | | |
|---|----------------------------|------------------|------------------|------------------------------|-----------------------------|----------------------------|------------------------------------|----------------|----------------------|----------------------------------|----------------|-------------|
| County | Birthweight ⁽⁵⁾ | | | Height and Weight | | | | | | Anemia Low Hb/Hct ⁽⁸⁾ | | |
| | Low | High | | Short Stature ⁽⁶⁾ | Under-weight ⁽⁷⁾ | Over-weight ⁽⁷⁾ | ≥ 2 Yrs Over-weight ⁽⁷⁾ | | | | | |
| | Number | % <2500 g (Rank) | % >4000 g (Rank) | Number | % <5th (Rank) | % <5th (Rank) | % >95th (Rank) | Number | % 85th- <95th (Rank) | % >95th (Rank) | Number | % (Rank) |
| Ulster | 1,091 | 9.1 (42) | 8.5 (24) | 5,120 | 3.6 (5) | 10.0 (61) | 11.4 (9) | 1,953 | 16.6 (20) | 18.2 (54) | 3,079 | 9.7 (12) |
| Warren | 579 | 9.0 (39) | 11.6 (53) | 4,066 | 6.0 (48) | 5.2 (29) | 12.0 (15) | 1,916 | 19.3 (55) | 14.9 (22) | 2,477 | 5.6 (1) |
| Washington | 615 | 8.5 (33) | 10.2 (45) | 4,537 | 6.0 (48) | 2.4 (5) | 16.2 (58) | 2,256 | 19.3 (55) | 16.8 (42) | 3,208 | 11.7 (26) |
| Wayne | 676 | 7.8 (25) | 8.9 (32) | 4,579 | 5.3 (34) | 4.5 (25) | 14.1 (36) | 2,136 | 19.3 (55) | 15.3 (28) | 3,148 | 17.0 (50) |
| Westcheste | 8,109 | 7.3 (17) | 7.5 (11) | 53,389 | 4.7 (23) | 5.1 (27) | 16.3 (59) | 24,655 | 17.2 (29) | 18.2 (54) | 35,502 | 13.7 (36) |
| Wyoming | 88 | * * | * * | 1,800 | 3.6 (5) | 4.1 (21) | 13.2 (28) | 815 | 17.1 (26) | 14.4 (16) | 1,198 | 9.5 (10) |
| Yates | 166 | 9.6 (53) | 13.3 (59) | 1,033 | 5.0 (27) | 3.3 (16) | 15.6 (54) | 481 | 20.0 (60) | 16.8 (42) | 748 | 12.7 (33) |
| New York | 167,993 | 8.2 | 7.6 | 1,285,221 | 4.2 | 7.2 | 12.8 | 568,518 | 16.5 | 16.3 | 863,791 | 13.1 |

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.