

**Table 6B**  
**2005 Pediatric Nutrition Surveillance** <sup>(1)</sup>  
**New York**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years** <sup>(2)</sup>

<b>Comparison of Growth and Anemia Indicators by County</b> <sup>(3,4)</sup>													
<b>County</b>	Birthweight <sup>(5)</sup>			Height and Weight							Anemia Low Hb/Hct <sup>(8)</sup>		
	Low	High		Short Stature <sup>(6)</sup>	Under- weight <sup>(7)</sup>	Over- weight <sup>(7)</sup>		≥ 2 Yrs Over- weight <sup>(7)</sup>					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% >95th (Rank)	Number	% (Rank)
Albany	603	7.6 (12)	7.5 (21)	4,600	4.0 (17)	7.4 (40)	10.9 (11)	2,124	17.8 (37)	12.7 (10)		3,064	19.7 (56)
Allegany	141	6.4 (6)	6.4 (14)	1,241	6.8 (57)	1.9 (2)	11.7 (26)	596	18.0 (41)	11.1 (4)		814	7.0 (6)
Bronx	7,171	10.9 (44)	6.2 (9)	56,758	3.5 (9)	8.0 (48)	13.5 (44)	26,253	15.8 (17)	17.7 (55)		39,830	13.6 (38)
Broome	517	7.4 (10)	11.2 (48)	4,350	6.1 (44)	4.2 (22)	13.9 (48)	2,105	17.2 (28)	13.4 (17)		2,983	7.5 (8)
Cattaraugu	207	5.3 (3)	10.1 (44)	1,913	6.3 (48)	2.1 (3)	13.6 (46)	903	18.3 (46)	14.7 (32)		1,321	8.9 (12)
Cayuga	174	9.8 (35)	6.9 (17)	1,632	4.8 (27)	7.4 (40)	12.8 (34)	808	15.8 (17)	13.1 (14)		978	14.9 (46)
Chautauqua	456	10.3 (38)	7.7 (24)	3,852	6.1 (44)	4.4 (23)	11.2 (17)	1,848	15.4 (12)	13.0 (13)		2,714	10.6 (24)
Chemung	291	10.0 (36)	6.2 (9)	2,623	6.1 (44)	3.2 (13)	11.3 (20)	1,322	19.0 (53)	12.4 (8)		1,849	7.4 (7)
Chenango	264	12.1 (49)	9.1 (40)	2,400	6.5 (53)	2.4 (5)	13.3 (41)	1,149	17.9 (39)	15.6 (38)		1,616	18.3 (51)
Clinton	186	7.0 (8)	8.1 (27)	1,461	5.3 (37)	10.0 (60)	8.8 (3)	659	18.1 (43)	12.4 (8)		984	5.9 (3)
Columbia	153	7.8 (16)	11.8 (51)	1,226	4.9 (30)	4.8 (26)	14.4 (51)	566	17.3 (29)	15.5 (35)		871	12.6 (35)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**2005 Pediatric Nutrition Surveillance** <sup>(1)</sup>  
**New York**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years** <sup>(2)</sup>

<b>Comparison of Growth and Anemia Indicators by County</b> <sup>(3,4)</sup>												
<b>County</b>	Birthweight <sup>(5)</sup>			Height and Weight							Anemia Low Hb/Hct <sup>(8)</sup>	
	Low	High		Short Stature <sup>(6)</sup>	Under- weight <sup>(7)</sup>	Over- weight <sup>(7)</sup>		≥ 2 Yrs Over- weight <sup>(7)</sup>				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% >95th (Rank)	Number
Cortland	170	7.1 (9)	8.8 (35)	1,320	6.4 (51)	2.7 (10)	11.3 (20)	596	15.1 (9)	12.1 (6)	866	9.8 (15)
Delaware	122	9.0 (28)	6.6 (15)	975	4.7 (25)	5.1 (27)	13.2 (40)	483	14.9 (7)	16.8 (47)	657	10.7 (25)
Dutchess	479	10.9 (44)	8.4 (33)	3,162	2.3 (1)	9.8 (58)	11.2 (17)	1,325	16.0 (20)	15.7 (40)	1,955	10.4 (21)
Erie	2,195	10.4 (40)	7.5 (21)	16,749	4.1 (19)	8.5 (51)	9.8 (5)	7,766	15.4 (12)	12.3 (7)	10,429	10.0 (17)
Essex	78	* *	* *	771	3.8 (11)	6.6 (35)	11.5 (23)	377	19.4 (56)	16.4 (44)	512	4.7 (2)
Franklin	151	8.6 (24)	6.0 (6)	1,302	5.8 (41)	3.5 (18)	11.2 (17)	621	17.4 (33)	14.8 (33)	894	11.3 (29)
Fulton	117	6.8 (7)	12.0 (52)	1,045	6.8 (57)	10.0 (60)	11.8 (27)	519	11.9 (1)	11.2 (5)	695	6.6 (4)
Genesee	225	8.9 (26)	5.8 (4)	1,827	5.1 (34)	6.3 (33)	14.1 (49)	867	17.9 (39)	16.8 (47)	1,209	14.6 (43)
Greene	80	* *	* *	593	4.2 (21)	4.6 (25)	12.8 (34)	284	16.2 (23)	13.7 (21)	424	12.3 (34)
Hamilton	14	* *	* *	93	* *	* *	* *	45	* *	* *	65	* *
Herkimer	156	7.7 (14)	10.9 (46)	1,474	7.1 (60)	3.1 (12)	12.5 (32)	683	15.2 (10)	13.9 (24)	1,013	29.0 (61)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**2005 Pediatric Nutrition Surveillance** <sup>(1)</sup>  
**New York**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years** <sup>(2)</sup>

<b>Comparison of Growth and Anemia Indicators by County</b> <sup>(3,4)</sup>												
<b>County</b>	Birthweight <sup>(5)</sup>			Height and Weight						Anemia Low Hb/Hct <sup>(8)</sup>		
	Low	High		Short Stature <sup>(6)</sup>	Under- weight <sup>(7)</sup>	Over- weight <sup>(7)</sup>		≥ 2 Yrs Over- weight <sup>(7)</sup>				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- <95th (Rank)			% >95th (Rank)
Jefferson	517	5.8 (4)	11.6 (50)	3,962	4.8 (27)	6.9 (37)	8.2 (1)	1,750	12.7 (2)	9.8 (1)	2,579	10.2 (19)
Kings	12,267	9.6 (33)	6.2 (9)	98,113	3.9 (13)	8.1 (49)	11.4 (22)	45,330	15.2 (10)	14.3 (30)	67,955	12.1 (32)
Lewis	158	9.5 (31)	7.0 (18)	1,297	4.2 (21)	7.3 (38)	10.3 (6)	632	14.9 (7)	13.8 (22)	868	14.2 (42)
Livingston	111	8.1 (18)	8.1 (27)	966	4.5 (24)	5.5 (29)	13.1 (39)	456	18.4 (47)	16.4 (44)	671	19.2 (55)
Madison	199	12.1 (49)	11.1 (47)	1,789	4.9 (30)	3.2 (13)	11.6 (24)	838	18.5 (50)	12.9 (12)	1,214	11.2 (28)
Monroe	1,507	10.3 (38)	7.0 (18)	12,057	3.3 (4)	9.5 (56)	11.0 (15)	5,617	16.1 (22)	14.1 (27)	7,775	10.5 (22)
Montgomery	127	8.7 (25)	7.9 (26)	1,268	5.7 (40)	6.5 (34)	14.9 (56)	623	15.6 (14)	16.4 (44)	876	14.6 (43)
Nassau	1,627	8.4 (21)	5.1 (2)	10,121	3.9 (13)	9.3 (55)	14.8 (54)	3,840	17.0 (26)	21.1 (59)	6,247	10.9 (26)
New York	6,328	8.9 (26)	6.2 (9)	48,404	3.0 (3)	7.7 (45)	12.0 (30)	21,649	17.4 (33)	15.6 (38)	33,812	11.1 (27)
Niagara	494	10.9 (44)	8.3 (31)	4,040	2.9 (2)	7.4 (40)	10.5 (7)	1,972	17.3 (29)	14.2 (29)	2,409	6.7 (5)
Oneida	586	9.7 (34)	8.5 (34)	5,148	5.8 (41)	6.0 (30)	10.9 (11)	2,405	16.9 (25)	13.6 (20)	3,557	23.4 (58)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**2005 Pediatric Nutrition Surveillance** <sup>(1)</sup>  
**New York**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years** <sup>(2)</sup>

<b>Comparison of Growth and Anemia Indicators by County</b> <sup>(3,4)</sup>												
<b>County</b>	Birthweight <sup>(5)</sup>			Height and Weight						Anemia Low Hb/Hct <sup>(8)</sup>		
	Low	High		Short Stature <sup>(6)</sup>	Under- weight <sup>(7)</sup>	Over- weight <sup>(7)</sup>		≥ 2 Yrs Over- weight <sup>(7)</sup>		Number      %  (Rank)		
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- <95th (Rank)			% >95th (Rank)
Onondaga	1,074	10.6 (41)	7.2 (20)	9,804	4.2 (21)	3.9 (21)	10.9 (11)	4,646	14.2 (6)	10.9 (3)	6,797	18.7 (54)
Ontario	204	7.4 (10)	13.2 (53)	1,658	4.7 (25)	3.5 (18)	13.8 (47)	755	18.0 (41)	15.4 (34)	1,147	16.3 (48)
Orange	929	9.3 (30)	7.6 (23)	6,821	5.4 (38)	8.1 (49)	10.5 (7)	3,053	15.6 (14)	13.4 (17)	4,730	12.8 (36)
Orleans	106	4.7 (2)	1.9 (1)	837	4.1 (19)	9.2 (53)	14.1 (49)	392	20.9 (61)	18.1 (57)	539	18.6 (52)
Oswego	387	7.8 (16)	8.3 (31)	3,464	6.5 (53)	2.6 (8)	14.7 (53)	1,678	18.4 (47)	17.0 (49)	2,423	9.9 (16)
Otsego	123	12.2 (51)	10.6 (45)	995	5.5 (39)	6.1 (31)	13.0 (37)	497	16.5 (24)	15.5 (35)	662	12.2 (33)
Putnam	76	* *	* *	527	6.8 (57)	6.8 (36)	17.8 (61)	203	13.8 (4)	24.6 (61)	320	15.0 (47)
Queens	7,002	8.5 (22)	6.3 (13)	47,642	3.6 (10)	9.2 (53)	11.8 (27)	19,179	16.0 (20)	17.0 (49)	31,078	9.2 (14)
Rensselaer	380	11.3 (47)	6.1 (7)	2,874	3.9 (13)	7.9 (46)	10.9 (11)	1,354	18.1 (43)	13.1 (14)	1,921	19.7 (56)
Richmond	906	10.7 (42)	5.8 (4)	6,749	3.3 (4)	7.6 (44)	12.8 (34)	2,899	17.3 (29)	17.1 (51)	4,369	8.7 (11)
Rockland	925	6.2 (5)	9.0 (37)	6,834	6.1 (44)	8.9 (52)	9.6 (4)	3,021	15.7 (16)	12.7 (10)	5,022	14.6 (43)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**2005 Pediatric Nutrition Surveillance** <sup>(1)</sup>  
**New York**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years** <sup>(2)</sup>

<b>Comparison of Growth and Anemia Indicators by County</b> <sup>(3,4)</sup>												
<b>County</b>	Birthweight <sup>(5)</sup>			Height and Weight						Anemia Low Hb/Hct <sup>(8)</sup>		
	Low	High		Short Stature <sup>(6)</sup>	Under- weight <sup>(7)</sup>	Over- weight <sup>(7)</sup>	≥ 2 Yrs Over- weight <sup>(7)</sup>					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- <95th (Rank)	% >95th (Rank)	Number	% (Rank)
Saratoga	230	12.2 (51)	9.1 (40)	1,751	5.1 (34)	7.9 (46)	10.7 (9)	804	18.4 (47)	13.3 (16)	1,180	16.3 (48)
Schenectad	380	12.4 (53)	6.1 (7)	2,863	3.4 (7)	7.5 (43)	12.3 (31)	1,307	15.8 (17)	13.4 (17)	1,966	24.2 (59)
Schoharie	61	* *	* *	644	4.0 (17)	4.5 (24)	16.0 (58)	328	18.9 (52)	15.9 (41)	453	7.9 (10)
Schuyler	42	* *	* *	403	7.9 (61)	3.2 (13)	14.9 (56)	188	19.1 (54)	17.6 (54)	281	7.5 (8)
Seneca	48	* *	* *	462	5.0 (33)	2.6 (8)	11.9 (29)	219	17.8 (37)	16.0 (42)	340	24.7 (60)
St. Lawren	161	8.1 (18)	5.6 (3)	1,459	6.4 (51)	3.5 (18)	13.3 (41)	698	13.8 (4)	16.0 (42)	1,010	17.2 (50)
Steuben	267	10.1 (37)	9.0 (37)	2,331	6.6 (55)	2.4 (5)	12.5 (32)	1,146	18.5 (50)	14.0 (25)	1,619	11.4 (31)
Suffolk	2,507	8.5 (22)	7.7 (24)	13,942	3.8 (11)	9.7 (57)	14.4 (51)	4,635	20.5 (59)	22.3 (60)	7,792	9.0 (13)
Sullivan	222	9.5 (31)	8.1 (27)	1,772	6.7 (56)	2.9 (11)	13.5 (44)	822	17.3 (29)	17.2 (52)	1,152	13.8 (40)
Tioga	143	7.7 (14)	11.2 (48)	1,372	5.2 (36)	2.5 (7)	13.3 (41)	680	20.6 (60)	14.6 (31)	917	13.7 (39)
Tompkins	184	7.6 (12)	8.2 (30)	1,508	3.9 (13)	6.2 (32)	8.6 (2)	717	13.1 (3)	10.3 (2)	1,019	13.8 (40)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

**Table 6B**  
**2005 Pediatric Nutrition Surveillance** (1)  
**New York**  
**Comparison of Growth and Anemia Indicators by County**  
**Children Aged < 5 Years** (2)

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under-weight (7)	Over-weight (7)	≥ 2 Yrs Over-weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% >95th (Rank)	Number	% 85th- <95th (Rank)	% >95th (Rank)	Number	% (Rank)
Ulster	272	10.7 (42)	9.2 (42)	1,716	3.3 (4)	9.9 (59)	11.6 (24)	669	17.0 (26)	17.5 (53)	1,040	10.3 (20)
Warren	190	11.6 (48)	8.9 (36)	1,389	6.3 (48)	7.3 (38)	10.7 (9)	637	19.6 (58)	13.8 (22)	862	2.2 (1)
Washington	144	9.0 (28)	9.0 (37)	1,562	5.8 (41)	2.2 (4)	17.2 (60)	770	19.2 (55)	17.7 (55)	1,128	10.1 (18)
Wayne	166	3.6 (1)	9.6 (43)	1,453	6.3 (48)	3.3 (16)	13.0 (37)	652	19.5 (57)	14.0 (25)	1,081	18.6 (52)
Westcheste	2,169	8.3 (20)	6.7 (16)	18,023	4.8 (27)	5.1 (27)	16.1 (59)	8,429	17.4 (33)	18.7 (58)	12,295	11.3 (29)
Wyoming	71	* *	* *	594	3.4 (7)	3.4 (17)	11.1 (16)	270	17.4 (33)	14.1 (27)	411	10.5 (22)
Yates	39	* *	* *	325	4.9 (30)	1.8 (1)	14.8 (54)	148	18.2 (45)	15.5 (35)	258	12.8 (36)
<b>New York</b>	<b>57,279</b>	<b>9.3</b>	<b>6.8</b>	<b>440,306</b>	<b>4.0</b>	<b>7.6</b>	<b>12.2</b>	<b>197,807</b>	<b>16.3</b>	<b>15.5</b>	<b>297,518</b>	<b>12.1</b>
<b>Nation (Prior Year)</b>	<b>1,633,363</b>	<b>9.3</b>	<b>6.8</b>	<b>6,600,078</b>	<b>6.4</b>	<b>4.7</b>	<b>14.1</b>	<b>2,822,660</b>	<b>16.1</b>	<b>14.8</b>	<b>4,020,684</b>	<b>13.7</b>

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

\* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.